

Covid-19 vaccination question guide

A.1	Nom de la personne à vacciner:			
A.2	Date of birth:			
A.3	Address:			
		Yes	No	Clinical implications
B.1	Have you ever had a confirmed Covid-19 infection?			Persons with confirmed Covid-19 infection should be vaccinated at least once. Please also consult your national guidelines.
	If yes, when did the symptoms start?	Date:		At the time of vaccination, the acute infection should have come to an end. Recommendations on the interval between acute infection and vaccination vary between 30 days and 6 months. Please also consult your national guidelines.
B.2	Have you ever been vaccinated against Covid-19?			If the patient has already received a full series of Covid-19 vaccinations depending on the vaccine, other patients should be prioritised.
	If yes, which vaccine did you get?	Date:	Vaccine:	
		Date :	Vaccine:	
B.3	Have you been vaccinated in the last 14 days against any other disease?			If yes, an interval of at least two to four weeks to the Covid-19 vaccination should be respected. Especially if the patient has been vaccinated with a live virus vaccine, e.g. against measles, an interval of four weeks is recommended. Please also consult your national guidelines.
B.4	Have you ever had an allergic reaction such as fainting, breathing problems, skin reactions or high fever after a former vaccination including the Covid-19 vaccination?			If yes, the patient should only be vaccinated if he or she carries an elevated risk of a Covid-19 infection <u>and</u> if an allergic shock can be treated adequately and immediately. The patient should be advised to stay for 30 minutes after vaccination.

		Yes	No	Clinical implications
C.1	Do you currently have an acute infection with fever such as a cold, flu or even Covid-19?			If yes, the vaccination should be postponed until the acute symptoms have come to an end.
C.2	Do you suffer from a chronic health condition such as diabetes, high blood pressure or cancer?			If yes, the patient should be treated as priority patient for vaccination against Covid-19.
C.3	Do you suffer from a disorder of the immune system or do you get a treatment that reduces the capacity of the immune system such as chemotherapy or drugs that suppress the immune system?			If yes, the patient should be treated as priority patient for vaccination against Covid-19. The patient should be informed that the efficiency of the vaccine may be reduced and an infection with Covid-19 may still occur. Therefore, the patient should definitely continue preventive behaviour.
C.4	Do you live with HIV or AIDS			If yes, the patient should be treated as priority patient for vaccination against Covid-19.
C.5	Do you suffer from a disorder of the blood coagulation like thrombosis?			If yes, the patient may carry an elevated risk of thrombosis if immunised with a vaccine of AstraZeneca, Johnson&Johnson or Sputnik. Then it is better to use an mRNA vaccine (Moderna, BionTech/Pfizer) if available.
C.6	Do you suffer from any other allergy?			If yes, the patient should only be vaccinated if he or she carries an elevated risk of a Covid-19 infection and if an allergic shock can be treated adequately and immediately. The patient should be advised to stay for 30 minutes after vaccination.
C.7	Are you currently pregnant?			If yes, the patient should be treated as priority patient for vaccination against Covid-19. All vaccines have been rated as safe for pregnant women in all trimesters.